arXiv:quant-ph/0112071v8 2 May 2002

# A simple way to detect the state transition caused by the nondiagonal abelian Berry phase

Wang Xiangbin*, Matsumoto Keiji†, Fan Heng‡ and Tomita Akihisa§

Imai Quantum Computation and Information project, ERATO, Japan Sci. and Tech. Corp.

Daini Hongo White Bldg. 201, 5-28-3, Hongo, Bunkyo, Tokyo 113-0033, Japan

Jian-Wei Pan**

Institute fur Experimentalphysik, Universitat Wein, Boltzmanngasse 5, 1090 Wien, Austria

## Abstract

In a nondegenerate system, the abelian Berry's phase will never cause transitions among the Hamiltonian's eigenstate. However, in a degenerate system, it is well known that the state transition can be caused by the non-abelian Berry phase. Actually, in a such a system, the phase factor is not always nonabelian. Even in the case that the phase is a nondiagonal matrix, it still can be an abelian phase and can cause the state transitions. We then propose a simple scheme to detect such an effect of the Berry phase to the degenerate states by using the optical entangled pair state. In this scheme, we need not control any external parameters during the quantum state evolution.

*email: wang@qci.jst.go.jp

†email: keiji@qci.jst.go.jp

‡email: fan@qci.jst.go.jp

§email: a-tomita@az.jp.nec.com

**email: pan@ap.univie.ac.at

Geometric phase [1–3] plays an important role in quantum interferometry and many other disciplines. Many observable quantum phyonemena are related to the geometric phase shift. The geometric phase to the degenerate states is qualitatively different from the one to the nondegenerate states, because it can cause the detectable effects which never happen to the nondegenerate states. For example, to the degenerate states one can obtain the non-abelian geometric phase [4–8]. Such a non-abelian Berry phase can cause transitions among the orthogonal states belonged the same eigenvalue of the Hamiltonian therefore could be useful in quantum computation [9]. Due to the non-abelian property, if we alternate two cyclic adiabatic operations, the states evolve differently. In this paper we will show that, to the degenerate states, there are *abelian* phase factors which are nondiagonal matrices. This is to say, even the Abelian Berry phase is qualitatively different from the one to the nondegenerate states, because it can cause the transitions among the eigenstates after a cyclic adiabatic evolution. We also give a feasible scheme to detect this type of nontrivil effect caused by the Berry phase to the degenerate states.

The experimental testing of geometric phase is an interesting and important topic. For a two level system, geometric phase is equal to half of the solid angle subtended by the area in the Bloch sphere enclosed by the closed evolution loop of the eigenstate. Most of the times, after certain time evolution path, there are both dynamic and geometric phase shift, this causes extra treatment to prove the effects of the geometric phase.

Recently, applications of the geometric phase is extended the geometric quantum computation [9,11–16], i.e., making the quantum gate through the geometric phase shift. In particular, several of them are based on the idea of using the Berry phase [9,14,15] shift to the degenerate states. Different from the case to the nondegenerate states, to a degenerate system, the Berry phase after a cyclic adiabatic evolution may cause the bit flip or the conditional bit flip between two states. This is rather significant because of its obvious fault tolerance property. For the nonabelian holonomic quantum computation, the computational space $C$ is always an eigen space(spanned by a number of degenerate eigen states). The first clear experimental proposal to produce the controllable Berry phase ship for the degenerate

system was raised by Duan et al [15]. The proposal suggests using trapped ions for the test. However, in this scheme, we will have to exactly control the external parameters so that the eigenvalues of the instantaneous Hamiltonian keeps to be zero. In this paper, we propose a simple scheme to test the states transition caused by Berry phase to the degenerate. Our scheme could be more robust because it requires little external parameters control in the whole process.

Lets first see what is the non-trivil effect caused by the arbitrary Berry phase to the degenerate states. Suppose orthonormal states $\{|\psi_i'\rangle\}$ spans a degenrate space of Hamiltonian $H$, i.e., all of the states are the eigenstates of $H$ with the same eigenvalue. If we change the Hamiltonian adiabatically and cyclically, there could be a non-trivil time evolution of the states in the degenerate subspace. There is no dynamical contribution to the time evolution because the dynamical phase is all zero. This time evlution operator in general can be a matrix with nonzero off diagonal elements. Since there is no dynamical contribution, we name this time evolution operator as the geometric phase factor. Suppose after certain cyclic adiabatic process the Berry phase factor is $U$, i.e.

$$\{|\psi_i'(t)|\rangle\} = U\{\psi_i'(0)\rangle\}. \tag{1}$$

Since $U$ is unitary, in principle we can always diagonalize it by another unitary matrix $S$ in the same degenerate space so that $S^\dagger US$ is diagonalized. This is to say, in the basis $\{|\psi_i\rangle\} = S\{|\psi_i'\rangle\}$, the time evolution operator is diagonal. Any state $|\psi_i\rangle = S|\psi_i'\rangle$ must be also an eigenstate of $H$ due to the degeneracy property. And, all states of $\{\psi_i\}$ have the same eigenvalues with that of $\{\psi_i'\}$. Consequently, if any adiabatical process causes a diagonal time evolution(a phase shift) in certain basis $\{|\psi_i\rangle\}$, the phase factor must be a non-diagonal matrix in another basis $\{|\psi_i'\}$. This type of time evolution is in general regarded as *nonabelian* and may cause transitions among the eigenstates, even though all states keep on being the eigenstates of of the instantaneous Hamiltonian during the whole process. This transition never happens by the same operation to the nondegenerate eigenstates. However, in certain case, the evolution operator can be *abelian* although it is not diagonal. And as we

will show in the following, such a nontrivil *abelian* phase can also cause state transitions.

Consider the following entangled state for the spin half system

$$|\Phi_{\pm}\rangle = \frac{1}{\sqrt{2}}(|0_A 1_B\rangle \pm |1_A 0_B\rangle), \tag{2}$$

subscripts $A$ and $B$ represent for the qubit $A$ or qubit $B$ respectively, 0 is spin down and 1 is spin up. Denoting $|\psi_1\rangle = |0_A 1_B\rangle$ and $|\psi_2 = |1_A 0_B\rangle$. Obviously state $|\psi_1\rangle$ and $|\psi_2\rangle$ are dengenrate in a $Z$-direction external field.

Now we consider how to produce an *abelian* and *nondiagonal* matrix phase factor to the two level degenerate states. We can first spatially separate spin $A$ from spin $B$. Obviously, we can also acquire a non-trivil Berry phase in the basis $|\Phi_{\pm}\rangle$ which is linearly superposed by state $|\psi_1\rangle$ and $|\psi_2\rangle$.

We will show that we can have a non-trivil effect if it is observed in the $|\Phi_{\pm}\rangle$ basis through the abelian phase in the basis $|\psi_{1,2}\rangle$. Consider another set of states $|\Phi_{\pm}\rangle$, each of which is a linear superposed state of $|\psi_1\rangle$ and $|\psi_2\rangle$. We can first spatially separate qubit $A$ and $B$. Then rotate the local field of qubit $B$ so that qubit $B$ evolve a *closed* loop. After the evolution, the Berry phase for $0_A$ and $1_A$ are opposite and the value is propotional to solid angle substended by the evolution loop at the degenerate point of the Bloch sphere for spin $B$. Explicitly, we have

$$\Gamma_{0,1} = \pm\Omega/2 \tag{3}$$

where $\Gamma_{0,1}$ is the Berry phase shift for state $|0_B\rangle$ or $|1_B\rangle$ and $\Omega$ is the value of solid angle substended by the evoultion loop. For example, if we rotate the field adiabatically around the $Z'$-axis for a loop. Then we have $|\Gamma_{0,1}| = \pi\cos\theta$, $\theta$ is the angle between $Z$ and $Z'$. Without loss of generality, we use $\Gamma$ to denote the Berry phase shift for state $|0_B\rangle$. At time $T$ when the evolution loop is completed we have

$$|\Phi(T)\rangle = \frac{1}{\sqrt{2}}(e^{-i\Gamma}|\psi_1\rangle \pm e^{i\Gamma}|\psi_2\rangle). \tag{4}$$

Note that here no dynamical phase is involved because state $|\psi_1\rangle$ and state $|\psi_2\rangle$ are degenerate in the whole process. Although the phase is diagonal in the $|\psi_{1,2}\rangle$ basis, it is *not*

diagonal in the $|\Phi_\pm\rangle$ basis. In such a non-trivil basis, the phase factor is given as

$$\Sigma = \begin{pmatrix} \cos\Gamma & -i\sin\Gamma \\ i\sin\Gamma & \cos\Gamma \end{pmatrix}. \tag{5}$$

This is to say, the time evolution is give as:

$$\begin{pmatrix} |\Phi_+(T)\rangle \\ |\Phi_-(T)\rangle \end{pmatrix} = \Sigma \begin{pmatrix} |\Phi_+(T)\rangle \\ |\Phi_-(T)\rangle \end{pmatrix}. \tag{6}$$

Note that there is no dynamical contribution to this no-commutative phase factor $\Sigma$ because in the whole adiabatic process we have

$$H(t)|\Phi_\pm(t)\rangle = 0. \tag{7}$$

The accumation of the dynamical phase is trivilly zero.

This matrix phase factor $\Sigma$ is an *abelian* phase factor although it can be non-diagonal. This is because all possible $\Sigma$ by the above operation commutate with each other, i.e., the matrices $\Sigma$ with different $\Gamma$ values commute. Although the phase factor here is abelian, it is different from the usual Berry phase in the nondegenerate which is diagonal and which never causes transitions among the eigenstates of the Hamiltonian.

Now we give a scheme to realize the above phase shift. It has been demonstrated extensively for the EPR state of eq(2) by the polarized photon pairs [17]. Now we are using polarized photons, state $|0\rangle$ and state $|1\rangle$ in eq(2) represent vertical or horizontal polarizotion respectively. A polarized photon evolves by the following time evolution operator when passing through a birefringence crystal [18]

$$U(s) = e^{-i\hat{\beta}\cdot\sigma\cdot s}. \tag{8}$$

Here $s$ is the the distance the light travels in the crystal, $\beta$ is the axis of the birefringence and $\sigma$ is the Pauli operator. Actually this is the spatial evolution operator. We can produce a non-abelian phase factor through the Berry phase shift for state $|\Phi_\pm\rangle$ in the following way:

1. Prepare the polarized entangled pair stae $|\Phi\rangle = |\Phi_+\rangle$ or $|\Phi\rangle = |\Phi_-\rangle$ and let the two photons are flying in the opposite way.

2. Let photon $A$ fly in the medium $A$ made of a crystal whose birefringence axis is along $Z$-axis. Meanwhile, photon $B$ is flying in the medium $B$ made up of the same crystal materials, however its birefringence axis changes gradually around $Z'$ axis for a loop. To meet the adiabatic condition we require the value $\mathrm{d}\hat{\beta}/ds$ very small. Mediums $A$ and medium $B$ have the same length. We set the length of the medium to be the value so that the refringence axis changes back to $Z$ direction.

3. After the two photons comes out the crystal, the state $|\Phi_{\pm}\rangle$ is changed to the one by eq(5).

Note that here the evolution path of photon $B$ on the Bloch sphere can be any irregular loop. The Berry phase is independent of the shape of the loop, it is only dependent on the area enclosed by the loop. In particular, if the birefringence $\hat{\beta}$ for medium $B$ is changed uniformly around $Z'$ axis and the angle between $Z$ and $Z'$ is set to $\theta = \pi/3$, the output state is $\Phi = \frac{1}{\sqrt{2}}(|\psi_1\rangle - |\psi_2\rangle)$, which is othorgonal to the initial state. To improve the efficiency, we can also let photon $A$ also fly in a medium whose birefringence axis gradually changes in a opposite manner as that of the medium $B$, i.e., polarization state of $A$ and $B$

In our schemes, we have used the EPR pair states. Obviously, we can also detect the non-abelian phase in the basis defined by the non-perfect entangled state, namely $|\Phi_+\rangle = \cos\beta e^{-i\alpha}|01\rangle + \sin\beta e^{i\alpha}|10\rangle$ and $|\Phi_-\rangle = \sin\beta e^{i\alpha}|01\rangle - \cos\beta e^{-i\alpha}|10\rangle$ where $\alpha$ and $\beta$ are non-zero real number satisfying the constraint $|\alpha|^2 + |\beta|^2 = 1$.

In summary, we have proposed simple schemes to produce and test the *abelian* but *nondiagonal* Berry phase. It should be interesting to take a further study on realizing the geometric controlled-NOT gate on the polarized photon qubits.

**Acknowledgement:** We thank Prof Imai for support. WXB thanks Prof N. Gisin, A. K. Pati and J. Pachos for fruitful discussions.

## REFERENCES


[1] S. Pancharatnam, Proc. India Acad. Sci. A **44,** 247(1956).

[2] M.V. Berry, Proc. R. Soc. London, Ser. A **392**, 45(1984).

[3] B. Simon, Phys. Rev. Lett., 51, 2167(1983).

[4] F. Wilczek and A. zee, Phys. Rev. Lett. 11, 2111(1984).

[5] C.A. Mead, Phys. Rev. Lett. 59, 161(1987); A. Zee, Phys. Rev. A38, 1(1988).

[6] J. W. Zwanziger, M. Koenig, and A. Pines, Phys. Rev. A42, 3107 (1990).

[7] J. Segert, J. Math. Phys., 28, 2102(1987); Ann. Phys. 179, 294(1987).

[8] J.E. Avron, L. Sadunl, J. Segert, and B. Simon, Phys. Rev. Lett., 61, 1329(1988); Commun. In Math. Phys.124, 595(1989).

[9] J. Pachos, P. Zanardi and M. Rasetti, Phys. Rev. A61, 010305(R)(2000).

[10] D. Suter, K. T. Muller, and A. Pines, *Phys. Rev. Lett.* **60**, 1216(1988).

[11] J A Jones, V Vedral, A Ekert, and G Castagnoli, *Nature* **403**, 869(2000)

[12] Falcl G, Fazio R, Palma G.M., Siewert J and Verdal V, *Nature* **407**, 355(2000)

[13] Xiaoguang Wang and P. Zanardi, quant-ph/0111017.

[14] P. Zanardi, M. Rasetti, Phys. Lett. A264, 94(1999)

[15] Duan L-M, Cirac J. I. and Zoller P, SCience 292, 1695(2001)

[16] Wang Xiang-bin and Matsumoto Keiji, Phys. Rev. Lett. 87, 097901(2001).

[17] D. Boumeester, et at, Nature, 390, 6660(1997)

[18] N. Gisin and A. Go, quant-ph/0004063